

Exercise 12.1 Derrick's Theorem

We have with $\phi'_i = \phi_i(\mathbf{x}/\alpha)$:

$$\begin{aligned} T' &= \frac{1}{2} \int d^D x \frac{1}{\alpha^2} \left(\nabla \phi_i \left(\frac{\mathbf{x}}{\alpha} \right) \right)^2 \\ &= \frac{1}{2} \int d^D (x\alpha) \alpha^{D-2} \left(\nabla \phi_i \left(\frac{\mathbf{x}}{\alpha} \right) \right)^2 \\ &= \alpha^{D-2} T \end{aligned}$$

and for the potential energy

$$U' = \int d^D x V \left(\phi_i \left(\frac{\mathbf{x}}{\alpha} \right) \right) = \alpha^D U.$$

We need $\alpha = 1$ to be a local minimum of the energy because this is nothing but the condition that the soliton should be stable against growing or shrinking.

$$\left. \frac{d}{d\alpha} E \right|_{\alpha=1} = (D-2)T + DU$$

which we set to zero to get

$$(2-D)T = DU$$

which is in contradiction to $T > 0$ and $U > 0$ for $D \geq 2$.

We can also argue with the sign of the derivative:

$$\frac{d}{d\alpha} E = (D-2)\alpha^{D-3}T + D\alpha^{D-1}U.$$

For $D \geq 2$ this is a positive function for all $\alpha > 0$, from this we conclude that the soliton spectrum extends down to $E = 0$ which is approached by spreading out the soliton more and more. Therefore by taking $\alpha \rightarrow 0$ we transform the soliton into the trivial vacuum ground state. This is in contradiction to the fact that ϕ_i was assumed to be a soliton solution.