

Phase Transitions and Critical Phenomena



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Exercise Sheet 3

HS 14

V. B. Geshkenbein

Problem 1 Two-component superconductivity in tetragonal crystals

Tetragonal crystals may exhibit superconductivity with a two component order parameter

$$\eta = (\eta_1, \eta_2). \quad (1)$$

The Landau free energy of such a superconducting order parameter is

$$F = \alpha \eta \cdot \eta^* + \frac{\beta_1}{2} (\eta \cdot \eta^*)^2 + \frac{\beta_2}{2} |\eta \cdot \eta|^2 + \frac{\beta_3}{2} (|\eta_1|^4 + |\eta_2|^4). \quad (2)$$

Find all possible superconducting phases as function of the coefficients $\beta_1, \beta_2, \beta_3$.

Problem 2 Susceptibility near the Néel temperature

Consider the Landau free energy of spins on a bipartite lattice with external field \mathbf{H}

$$F = \frac{a}{2} (M_1^2 + M_2^2) + A \mathbf{M}_1 \cdot \mathbf{M}_2 + \frac{b}{4} (M_1^4 + M_2^4) - \mathbf{H} \cdot \mathbf{M} \quad (3)$$

where $\mathbf{M}_1, \mathbf{M}_2$ are magnetizations of spins on the two sublattices and $\mathbf{M} = \mathbf{M}_1 + \mathbf{M}_2$ is their sum.

We argued in the lecture that close to the Néel temperature T_N

$$A = a - \alpha (T - T_N) \quad (4)$$

Find the magnetic susceptibilities $\chi_{\parallel}, \chi_{\perp}$ of the system near the Néel temperature for a field oriented parallel and perpendicular to the order parameter.