Phase Transitions and Critical Phenomena

ETH	Exercise Sheet 13	HS 14
Eidgenössische Technische Hochschule Zürich		V B Cochlephoin
Swiss Federal Institute of Technology Zurich		V. D. Gesnkenbein

Problem 1 Vertex renormalization of the *n*-component model

Consider again the n-component model from the lecture with a quartic term

$$u \int d^d \boldsymbol{r} F_{ijk\ell} \phi_i(\boldsymbol{r}) \phi_j(\boldsymbol{r}) \phi_k(\boldsymbol{r}) \phi_\ell(\boldsymbol{r})$$
(1)

where $F_{ijk\ell}$ is the completely symmetric tensor of rank four,

$$F_{ijk\ell} = \frac{1}{3} \left(\delta_{ij} \delta_{k\ell} + \delta_{ik} \delta_{j\ell} + \delta_{i\ell} \delta_{j,k} \right).$$
⁽²⁾

Prove the identity

- -

$$\sum_{k\ell} \left(F_{abk\ell} F_{k\ell cd} + F_{ack\ell} F_{k\ell bd} + F_{adk\ell} F_{k\ell bc} \right) = \frac{n+8}{3} F_{abcd} \tag{3}$$

that is needed to find the RG equation for the quartic coefficient u.

Problem 2 ε -expansion of the *n*-component model

Use identity (3) for the vertex renormalization and identity

$$\sum_{k} F_{ijkk} = \frac{n+2}{3} \delta_{ij} \tag{4}$$

for the propagator renormalization to arrive at the following RG equations for the n-component model,

$$\frac{du}{d\xi} = -\frac{n+8}{9}u^2\tag{5}$$

$$\frac{d\tau}{d\xi} = -\frac{n+2}{9}u\tau\tag{6}$$

Use these to derive the critical exponent γ in $4 - \varepsilon$ dimensions to the linear order in ε .